

#904803504

ID 88655301

QE
98
V74
1981

Geology and Mineral Resources
of the

Cache Creek-Rocky Creek
Wilderness Study Area

By

Larry Vredenburg
August, 1981

BLM Library
Bldg. 50
Denver Federal Center
P.O. Box 25047
Denver, Colorado 80225

RECEIVED
JAN 4 7 57 AM '81

SUMMARY

The Cache Creek-Rocky Creek WSA contains deposits or occurrences of mercury-gold, asbestos, placer chromite, gemstones, magnesite and sand and gravel. There has been production of mercury from within the area and there is potential for production of gold. Recent development work at the Knoxville mercury mines has delineated a major gold deposit which may extend into the WSA. Claims have been staked in the WSA to explore for a possible extension of the gold mineralization at Knoxville.

Natural gas seeps occur within the WSA, and there is potential for oil and gas. There is potential for the production of electrical energy from geothermal sources at Wilbur Springs and the geothermal reservoir may extend into the WSA.

The first of these is the fact that the amount of energy which is converted into heat in the process of the combustion of a fuel is not constant, but varies with the conditions of the combustion. This is due to the fact that the rate at which the fuel is consumed is not constant, but varies with the conditions of the combustion. The second of these is the fact that the amount of energy which is converted into heat in the process of the combustion of a fuel is not constant, but varies with the conditions of the combustion. This is due to the fact that the rate at which the fuel is consumed is not constant, but varies with the conditions of the combustion.

The third of these is the fact that the amount of energy which is converted into heat in the process of the combustion of a fuel is not constant, but varies with the conditions of the combustion. This is due to the fact that the rate at which the fuel is consumed is not constant, but varies with the conditions of the combustion. The fourth of these is the fact that the amount of energy which is converted into heat in the process of the combustion of a fuel is not constant, but varies with the conditions of the combustion. This is due to the fact that the rate at which the fuel is consumed is not constant, but varies with the conditions of the combustion.

INTRODUCTION

The Cache Creek-Rocky Creek Wilderness Study Area (WSA) is located in the Coast Range of California, east of Clear Lake, south of Wilbur Spring, and north of Knoxville (see Figure 1). It encompasses 33,582 acres and embraces Buck Mountain, Round Mountain, Little Blue Ridge and Brushy Sky High. Portions of T.13N., R.5 and 6W. and T.12N., R.5W.MDM are within this area. It is located within the Clear Lake Resource Area of the Ukiah District.

Access into the WSA is restricted to a jeep trail from Highway 20 to Cache Creek across the central part of the area. During a fire in the summer 1981, a fire break was bulldozed from Bear Spring to the north for a distance of about two miles.

PREVIOUS WORKS

The first workers to map and describe the area were Becker (1888) and Forstner (1903) who studied the mercury resources of the area. Lawton (1956) mapped and described in detail the geology of the eastern portion of the Cache Creek-Rocky Creek WSA. A condensation of James Brice's PhD thesis of 1950 from the University of California, Berkeley, was published in 1953 by the California Division of Mines. The published thesis is a reconnaissance of the western part of the WSA. Other workers in the area include Moisseeff (1966), who mapped and described the Wilbur Springs district; Averett (1945), who described and mapped the Knoxville district; and Christine Carlson (1981), who recently mapped and described an area south and east of Wilbur Springs.

In 1972 the U.S. Geological Survey began in-depth studies of the Geysers geothermal field. The results of these studies were published in McLaughlin and Donnelly-Nolan (1981).

SCOPE OF REPORT

This report was written to identify the mineral resources of the Cache Creek-Rocky Creek WSA. It was written entirely from published and unpublished literature sources and from personal and written communication with persons familiar with the area. No time was spent in the field. Because of the recent gold discoveries at Knoxville and the presence of a mercury-gold deposit deep within the WSA, a section discussing various aspects of this type of ore deposit has been included.

INTRODUCTION

The Great Smoky Mountains National Park (GSMNP) is located in the southeastern corner of the United States, east of the Appalachian Mountains. It is a large area of land, covering 520,000 acres, and is one of the largest national parks in the United States. The park is located in the southeastern corner of the United States, east of the Appalachian Mountains. It is a large area of land, covering 520,000 acres, and is one of the largest national parks in the United States.

The park is located in the southeastern corner of the United States, east of the Appalachian Mountains. It is a large area of land, covering 520,000 acres, and is one of the largest national parks in the United States. The park is located in the southeastern corner of the United States, east of the Appalachian Mountains. It is a large area of land, covering 520,000 acres, and is one of the largest national parks in the United States.

THE PARK

The park is located in the southeastern corner of the United States, east of the Appalachian Mountains. It is a large area of land, covering 520,000 acres, and is one of the largest national parks in the United States. The park is located in the southeastern corner of the United States, east of the Appalachian Mountains. It is a large area of land, covering 520,000 acres, and is one of the largest national parks in the United States.

The park is located in the southeastern corner of the United States, east of the Appalachian Mountains. It is a large area of land, covering 520,000 acres, and is one of the largest national parks in the United States. The park is located in the southeastern corner of the United States, east of the Appalachian Mountains. It is a large area of land, covering 520,000 acres, and is one of the largest national parks in the United States.

THE PARK

The park is located in the southeastern corner of the United States, east of the Appalachian Mountains. It is a large area of land, covering 520,000 acres, and is one of the largest national parks in the United States. The park is located in the southeastern corner of the United States, east of the Appalachian Mountains. It is a large area of land, covering 520,000 acres, and is one of the largest national parks in the United States.




Figure 1

Index Map - showing location of gold bearing mercury deposits.

RELIEF AND DRAINAGE

The highest elevation in the area is the 3,196 foot high Brushy Sky High. The lowest is at about 800 feet, located on Cache Creek. The ridges are generally formed by units of sandstones or serpentine which are more resistant to erosion than the adjacent shales in this area.

Cache Creek and its tributaries drain an area of about 750 square miles and flow through the WSA. Almost all of the streams in the area are tributary to Cache Creek. Cache Creek appears to be a superimposed stream whose intrenched meanders were formed on an earlier preexistent flood plain.

ACKNOWLEDGEMENTS

I would like to thank the following persons who helped in the preparation of this report: Susan Skinner, BLM Ukiah District Office; Edmond Lawrence, consultant; Christine Carlson, GS Menlo Park; Robert McLaughlin, GS Menlo Park; Don White, GS Menlo Park; M.C. Smith, Claimant; Steve Graham, Stanford University; Foster Howland, Resource Exploration.

GEOLOGY

According to McLaughlin (1981):

The northern California Coast ranges east of the San Andreas Fault...consist mainly of two approximately coeval units now separated by a great thrust (Bailey and others, 1970). The upper plate of the Coast Range thrust consists of a fragmented ophiolite complex of Late Jurassic Age (the Coast Range ophiolite), considered to represent oceanic crust (Bailey and others, 1970; McLaughlin and Pessagno, 1978) overlain depositionally by moderately deformed marine sedimentary rocks referred to as the Great Valley Sequence (Bailey and others, 1964). The Great Valley Sequence ranges in age from Late Jurassic to Late Cretaceous and consists of coarse ophiolite breccia or tuff near the base (McLaughlin and Pessagno, 1978) overlain by conglomerate, mudstone, and sandstone... The basal part of the Great Valley Sequence was largely derived from the depositionally underlying Coast Range ophiolite (McLaughlin and Pessagno, 1978).

Rocks in the lower plate of the Coast Range thrust have been assigned to the Franciscan assemblage (or Franciscan Complex of Berkland and others, 1972) and consist of a heterogeneous assemblage of intensely deformed and mildly to moderately metamorphosed sandstone, shale, chert, and mafic igneous rocks. Serpentinite, limestone, amphibolite, eclogite and high grade blueschists are minor but significant constituents. Franciscan rocks and their equivalents are now known to extend along the

Pacific coast of North America at least from Baja California, Mexico, to southern Alaska (Jones and others, 1977). Initial deformation and metamorphism of these rocks apparently occurred in Cretaceous and early Tertiary time, as the result of oblique northeast-directed subduction and strike-slip.

The Great Valley Sequence which is extensively exposed in the WSA consists of about 40,000 feet of Jurassic and Cretaceous shale, siltstone, sandstone, and occasional lenses of conglomerate and limestone. The sequence has been divided into twelve formations (see Figure 2). In the southern part of the WSA at the north end of Morgan Valley, the Knoxville Formation of the Great Valley Sequence is in depositional contact with serpentine (Lawton, 1956).

Much of the mercury and all of the magnesite and asbestos in the Coast Ranges occurs in altered serpentine. Hot solutions rich in SiO_2 and CO_2 rising along faults have replaced serpentine by silica-carbonate rock, which consists of chalcedony, opal, quartz, magnesite and calcite.

There are three areas within the WSA where the Franciscan Complex is exposed, as shown on the 1:250,000 scale state geologic map. One area is near Wilbur Springs where it is in fault contact with serpentine. The other two are in Deadman Canyon. The contacts with serpentine are shown as depositional in Deadman Canyon, but the exact correlation of the sediments is in doubt, and they may actually be Great Valley sediments.

The Cache Formation of Pliocene age is located in the northeast part of the WSA. This formation consists of a sequence of lacustrine clay, silt, sand and gravel beds with minor amounts of tuff that are from 1,000 to 6,500 feet thick. Identified terrestrial vertebrate fossils recovered from this unit, suggest an age range of about 1.8 to 3.0 million years (Casteel and Rymer, 1981).

The Quaternary age Clear Lake Volcanics range in age from 2 million years (m.y.) to about 10,000 years and range in composition from basalt to rhyolite. None of these rock units are exposed within the WSA. Basalt does occur nearby at Quackenbush Mountain, near the junction of Ferris Canyon and Cache Creek, at the Knoxville mines, and on Coyote Peak near Wilbur Springs.

PALEONTOLOGY

The Cache Creek-Rocky Creek WSA contains a great diversity of vertebrate and invertebrate fossils. Lawton's (1956) thesis covering the north half of the Morgan Valley quadrangle and the south half of the Wilbur Springs quadrangle carefully details the fossils present in each formation. Casteel and Rymer's (1981) papers describe several fossil fish discovered in the Cache Formation. The reader is referred to these works for further details.

Pacific coast of North America at least from Baja California
Mexico, to western Alaska (Jones and others, 1937). Initial
deformation and metamorphism of these rocks apparently occurred
in Late Proterozoic and early Paleozoic time, as the result of orogenic
movements directed northeastward and strike-slip.

The Great Valley Sequence which is extensively exposed in the KSA consists
of about 40,000 feet of Jurassic and Cretaceous shales, siltstones, sand-
stones, and occasional lenses of conglomerate and limestone. The sequence
has been divided into twelve formations (see Figure 2). In the southern
part of the KSA at the north end of Whipple Valley, the Knoxville Formation
of the Great Valley Sequence is in depositional contact with the
Knoxville (1937).

Most of the country and all of the magmatic and sedimentary in the Coast
Range occur in a single magmatic unit. The unit is 2500 and
3000 feet thick. It is a single unit, as indicated by a single
unit, which consists of a variety of quartz, quartzite, and calcite.

There are three areas within the KSA where the Franciscan Complex is ex-
posed, as shown on the 1:250,000 scale geologic map. One area is
near the mouth of the KSA where it is in contact with the Franciscan Complex. The
other two are in the Coast Range. The contacts with the Franciscan Complex are
not well defined in the Coast Range, but the correlation of the sedi-
ments is in doubt, and they are actually the Great Valley Sequence.

The Knox Formation of Whipple Valley is located in the northeast part of
the KSA. This formation consists of a sequence of sandstone, siltstone, and
shale, and is about 1000 feet thick. It is in contact with the Knox
Formation of Whipple Valley, which is about 1000 feet thick. The Knox
Formation of Whipple Valley is in contact with the Knox Formation of Whipple
Valley, which is about 1000 feet thick. The Knox Formation of Whipple
Valley is in contact with the Knox Formation of Whipple Valley, which is
about 1000 feet thick.

The Great Valley Sequence is a sequence of sedimentary rocks in the KSA. It
is about 40,000 feet thick and is composed of sandstone, siltstone, and
shale. It is in contact with the Knox Formation of Whipple Valley, which
is about 1000 feet thick. The Knox Formation of Whipple Valley is in
contact with the Knox Formation of Whipple Valley, which is about 1000
feet thick. The Knox Formation of Whipple Valley is in contact with the
Knox Formation of Whipple Valley, which is about 1000 feet thick.

REFERENCES

The Knox Formation of Whipple Valley is a sequence of sedimentary rocks in
the KSA. It is about 1000 feet thick and is composed of sandstone, silt-
stone, and shale. It is in contact with the Knox Formation of Whipple
Valley, which is about 1000 feet thick. The Knox Formation of Whipple
Valley is in contact with the Knox Formation of Whipple Valley, which is
about 1000 feet thick. The Knox Formation of Whipple Valley is in
contact with the Knox Formation of Whipple Valley, which is about 1000
feet thick. The Knox Formation of Whipple Valley is in contact with the
Knox Formation of Whipple Valley, which is about 1000 feet thick.





Figure 2

Northern Great Valley Sequence: composite column, Cache Creek to Wilbur Springs (from Page, 1966).

STRUCTURE AND TECTONICS

Faulting

Most of the folding and faulting (see Figure 3) in the area trends northwest-southeast. Some faults strike between north-south and north-northeast-south southwest. There is evidence that the Bear Valley and Hunting Faults were active before Late Cretaceous time. The Kennedy, Wilson and Hunting Faults have been active since Pliocene-Pleistocene time.

Folding

Folds have been mapped in four areas within the WSA. These are Grizzly Canyon northeast of Grizzly Spring, north of Round Mountain from Brushy Creek to Jack Canyon, at Dunfield flats and northwest of Morgan Valley.

Along Grizzly Canyon is a complex structure named by Lawton (1956) as the Grizzly structure. The structure consists of three elongate masses of serpentine with associated shale and sandstone, all of which have pierced sediments of the Knoxville Formation. Folds associated with the piercement are ill-defined. Beds in the surrounding sediments dip gently away from the piercement mass.

North of Round Mountain is a syncline named by Lawton (1956) as the central syncline. The axis of the syncline is sinuous, but trends generally east-west and plunges to the west. On the east, the axis of the syncline aligns with a zone of complex structures. Dunfield Flats is the western end of the zone of complex structures. In this zone, there are small tight folds and faults of apparently small displacement. North of Morgan Valley there are several west-northwest trending and southeast plunging parallel folds.

Siesmicity

No epicenters of earthquakes 4.0 on the Richter Scale or larger have been recorded in the WSA since 1900 (Real and others, 1978). Since 1969, when recording equipment was established in the area, there have been three earthquakes with magnitudes of less than 2 centered in the western part of the WSA. Between Morgan Valley and Lake Berryessa, there have been numerous earthquakes. The largest magnitudes are between 2.0 and 3.0. At Knoxville, there have been five earthquakes with magnitudes less than 2.0. The seismicity of the Geysers-Clear Lake area, is discussed by Bufe and others (1981).

GEOLOGIC HISTORY

During late Jurassic through Late Cretaceous time, sediments of the Franciscan Complex and the Great Valley Sequence were simultaneously being deposited in parallel basins on oceanic crust. The rocks of the Franciscan Complex are interpreted to have been deposited in a trench over an east-dipping subduction zone located to the west of the fore-arc basin of

Faulting

Most of the faulting and faulting (see Figure 2) in the area trends northwest-southeast. Some faults strike between north-south and north-northeast-south-southwest. There is evidence that the Bear Valley and Harding faults were active before late Pleistocene time. The Harding fault and Harding fault have been active since Pleistocene-Pleistocene time.

Faulting

There have been mapped in four areas within the WSA. There are mainly faults north-south of Harding fault, north of Round Mountain from Bear Valley to Lake Canyon, at Lakefield Lake and north-south of Morgan Valley.

Along Harding Canyon is a complex structure noted by Jackson (1955) as the Harding fault. The structure consists of three elongate masses of sedimentary rock separated by faults and separated by faults. This association with the Harding fault is the Harding fault. Beds in the surrounding sedimentary are mostly away from the Harding fault.


North of Round Mountain is a syncline noted by Jackson (1955) as the Harding fault. The axis of the syncline is horizontal, but trends generally east-west and widens to the west. In the east, the axis of the syncline aligns with a line of complex structures. Harding fault is the western end of the line of complex structures. In this area, there are small faults and faults of sporadically small displacements. North of Morgan Valley there are several west-northwest trending and southeast plunging faults.

Stratigraphy

An outcrop of sedimentary rock on the Harding fault at Lakefield has been mapped in the WSA since 1955 (see Figure 2). Since 1955, when the Harding fault was established in the area, there have been three outcrops with magnitudes of less than 2 centimeters in the western part of the WSA. Between Morgan Valley and Lake Canyon, there have been three outcrops. The largest outcrops are between 2.0 and 3.0 ft. However, there have been two outcrops with magnitudes less than 2.0 ft. The stratigraphy of the Harding-Clear Lake area is discussed by this author (1955).

Geologic History

During late Pleistocene through late Pleistocene time, sediments of the Harding Canyon and the Harding Valley sediments were substantially being deposited in parallel basins on opposite sides. The rocks of the Harding Canyon were interpreted to have been deposited in a trench over an east-dipping subduction zone located to the west of the Harding basin of




GEOLOGIC STRUCTURE MAP
OF MORGAN-WILBUR AREA

SCALE

1 1/2 0 1 2 miles

- Fault trace
- - - Inferred fault trace
- - - Permeability contact
- + Anticlinal axis
- + Synclinal axis

Figure 3



GEOLOGICAL STRUCTURE AND
OF MORGAN-WELLS AREA

1948

Scale 1:25,000

Figure 2

the Great Valley Sequence. However, several lines of evidence presented by McLaughlin (1981) suggests this may be an over-simplification of the sequence of events. The Great Valley Sequence is interpreted to represent fore-arc basin deposits that were derived from the Klamath and Sierran Island arc terraines as a series of coalescing submarine fans. The source areas for the Franciscan sediments have been destroyed by subduction.

From Mid-Cretaceous time onward, the rocks of the Great Valley Sequence and Franciscan Complex were simultaneously deformed by underthrusting caused by sea floor spreading. Superposition of the two units was brought about by the thrust faults, whose position today is nearly everywhere marked by serpentine. Beneath the thrust, the Franciscan rocks are sheared and metamorphosed.

During Tertiary time shearing along the San Andreas fault system has produced a zone at least 60 Km wide, which was affected by accumulation of strain and creep on numerous subsidiary fault zones. These faults, particularly those which trend northwest and north-northwest have provided major conduits for movement of thermal fluids to the surface (Hearn and others, 1981).

Following the deposition of the Cache Formation, the Coast Ranges experienced considerable uplift and deformation. Prior to the uplift, Cache Creek was contained in a wide valley or flood plain, which may have been at about the level of the Sacramento Valley during Pliocene and Pleistocene time (Lawton, 1956).

MINERAL RESOURCES

INTRODUCTION

Locatable minerals within the Wilderness Study Area (WSA) include mercury, gold, asbestos and chromite. In addition, adjacent to the area at Wilbur Springs to the north, at Knoxville to the south, at the Baker Mine to the southwest, and at Sulphur Bank to the west, gold-mercury ore has been reported (Becker, 1888). Mercury has been the primary metal sought at these mines, although gold was mined for several decades in the 1880s at Wilbur Springs. Recent development work at the Knoxville mines has delineated a major gold deposit which may extend into the WSA. Claims have been staked in the WSA to explore for this possible extension of the gold mineralization.

Oil, gas, and geothermal energy are leasable resources which potentially could occur in economic quantities. There are gas seeps within the WSA, and there are numerous oil and gas seeps at Wilbur Springs, and oil seeps are reported at Knoxville (Lawton, 1956, Averitt, 1945). There are hot springs at Wilbur Springs and Knoxville. At Wilbur Springs, it is believed that electric power could be generated. The hot water reservoir at Wilbur Springs may extend into the WSA.

The Great Valley Sequence. However, several lines of evidence presented by Wright (1961) suggest this may be an oversimplification of the sequence of events. The Great Valley Sequence is interpreted to represent the basin deposits that were derived from the Klamath and Sierra Nevada and deposited as a series of overlapping successions. The source areas for the Franciscan sediments have been destroyed by subduction.

From the Franciscan line south, the rocks of the Great Valley Sequence and Franciscan Complex were simultaneously deformed by unroofing caused by the Great Valley Sequence. Repositioning of the two units was brought about by the Great Valley Sequence, whose position today is nearly everywhere defined by a massive thrust. Beneath the thrust, the Franciscan rocks are located and metamorphosed.

During Tertiary time shearing along the San Andreas fault system has produced a zone at least 40 km wide, which was affected by accumulation of strain and stress on numerous subsidiary fault zones. These faults, particularly those which trend northwest and north-northeast have provided major conduits for movement of material fluids to the surface (Heintz and others, 1960).

Following the deposition of the Franciscan Formation, the Great Valley Sequence was deformed by uplift and deformation. Prior to the Tertiary, Great Valley Sequence was contained in a wide valley or flood plain, which may have been at about the level of the Sacramento Valley during Miocene and Pliocene time (Wright, 1962).

MINERAL RESOURCES

INTRODUCTION

Extensive studies within the Klamath Study Area (KSA) indicate extensive gold, silver and copper resources. In addition, adjacent to the area are lithium, boron, and other minerals. The Klamath Study Area is situated in the north, of Klamath County, at the Klamath River in the north, and at the Klamath River in the south. Gold-mining has been reported (Wright, 1962). However, the present study is limited to the Klamath River, although gold has been reported in the Klamath River in the 1880s at Klamath Falls. Recent development work at the Klamath River has indicated a major gold resource which may extend into the KSA. Claims have been filed in the KSA to explore for this possible extension of the gold resource.

Oil, gas, and geothermal energy are valuable resources which potentially exist in the Klamath Study Area. There are gas seeps within the KSA, and there are numerous oil and gas seeps at Klamath Falls and all seeps are reported at Klamath Falls (Wright, 1962). There are also seeps at Klamath Falls and Klamath River. At Klamath Falls, it is believed that geothermal energy could be generated. The hot water reservoir at Klamath Falls may extend into the KSA.

HISTORY

Beginning in the early 1860s, discovery and subsequent development of mineral resources took place at Wilbur Springs and Knoxville. Mercury was discovered at Knoxville in 1862 while cutting a grade for a road. The mines were important producers until 1883. Production slacked, but mercury was produced almost every year since the opening of the mines until 1943 (the last year figures are available). Table 1 lists production from the mines in this area. Past production of mercury from the Knoxville mines is significant. Knoxville ranks third in the United States in terms of total mercury produced.

At Wilbur Springs, mercury was discovered in 1862 and production took place at the Abbott Mine between 1870 and 1890, and at other periods of high mercury prices. Gold was discovered at the Manzanita Mine in the early 1860s and reportedly was worked principally for gold until 1891, when mercury was also mined. Widely known oil and gas seeps near Wilbur Springs were the object of unsuccessful drilling in 1865-1866 (Carlson, 1962, p. 23).

MERCURY-GOLD MINERALIZATION

The recent discovery of a large, previously unsuspected quantity of gold at Knoxville (at last estimate 3.2 million ounces), is not the only known occurrence of gold with mercury (or antimony) in the Coast Range of California. The earliest and perhaps one of the most comprehensive studies of the mercury deposits of the Pacific Coast was that of Becker (1888). He determined the presence of gold at mercury mines at Wilbur Hot Springs, Sulphur Bank, as well as the Knoxville mines. He also reported the occurrence of gold at the Picacho Mine, located southwest of New Idria. Other workers (Bradley 1916, Newmont Exploration 1965, Bailey and Meyers 1949) have identified gold at the Calistoga, Oat Hill mines, and the Stayton District.

Although over 50 percent of the largest mercury deposits in the Coast Range are associated with altered serpentine, mercury also occurs in rocks of the Franciscan and Knoxville Formations, as well as Tertiary volcanic rocks and at active hot springs. The rock type does not appear to be the controlling factor in the deposition of mercury in this area.

The presence of young high-angle faults appears to be the most important factor in the location of mercury deposits. All mercury deposits are associated with faults and fracture zones in this area. Averitt (1945) suggested that clay gouge zones effectively channeled mineral bearing water and this controlled deposition at the Knoxville mine.

The origin of the mercury, gold, hydrocarbons and the water, which acted and still acts as a transporting medium, are all intimately associated with the marine sediments. In a study of thermal and mineral waters in

Beginning in the early 1900s, discovery and subsequent development of mineral resources took place at Wilbur Springs and Knoxville. Wilbur was discovered at Knoxville in 1902 while cutting a grade for a road. The mine was important producer until 1907. Production slowed, but was not abandoned until every year since the opening of the mine until 1907 (the last year figures are available). Table 1 lists production from the mine in this area. Great production of mercury from the Knoxville mines is significant. Knoxville ranks third in the United States in terms of total mercury produced.

At Wilbur Springs, mercury was discovered in 1902 and production took place at the Wilbur mine between 1902 and 1907, and at other periods of this century period. Gold was discovered at the Knoxville mine in the early 1900s and reportedly was worked principally for gold until 1907. Wilbur was also mined. Wilbur was an oil and gas seep near Wilbur Springs were the object of unsuccessful drilling in 1902-1904 (Carson, 1907, p. 131).

MINERAL-TOO MINERALIZATION

The mineral discovery of a large, previously unexplored quantity of gold at Knoxville (at last estimate 2.5 million ounces), is not the only known occurrence of gold with mercury for enrichment in the Coast Range of California. The earliest and perhaps one of the most comprehensive studies of the mercury deposits of the Pacific Coast was that of Becker (1908). He determined the occurrence of gold at mercury sites at Wilbur Springs, Wilbur, and as well as the Knoxville mine. He also reported the occurrence of gold at the Placerville mine, located east of New York. Other reports (Baker 1910, Hanson, Exploration 1910, Baker and Meyer 1910) have indicated gold at the California, New York mine, and the Oregon mine.

Historically over 30 percent of the largest mercury deposits in the Coast Range are associated with active volcanoes, mercury also occurs in rocks of the Franciscan and Knoxville formations, as well as tertiary volcanic rocks and active hot springs. The rock type does not appear to be the controlling factor in the deposition of mercury in this area.

The presence of young high-angle faults appears to be the most important factor in the location of mercury deposits. All mercury deposits are associated with faults and fracture zones in this area. Austin (1962) suggested that clay gouge zones effectively channel (store) bearing water and this controlled deposition at the Knoxville mine.

The origin of the mercury, gold, hydrocarbons and the water, which acted and still acts as a transporting medium, are still unknownly associated with the entire continent. In a study of mineral and hydrothermal waters in

Year	Knoxville (Excelsior, Redington, Boston)	Reed (California, Andalusia)	Manhattan (Lake)	Red Elephant	Harrison (New England, Ruby, Grizzly, January)	Total ¹	Average price of quicksilver per flask
1860	---	---	---	---	---	---	53.55
61	---	---	---	---	---	---	42.05
62	444	---	---	---	---	444	36.35
63	852	---	(a)	---	---	852	42.08
64	1,914	---	(a)	---	---	2,714	45.90
65	3,545	---	(a)	---	---	3,545	45.90
66	2,254	---	(a)	---	---	2,254	53.13
67	7,862	---	(a)	---	---	7,862	45.90
68	8,686	---	(a)	---	---	9,808	45.90
69	5,018	---	(a)	---	---	6,598	45.90
1870	4,546	(b)	(a)	---	---	6,766	57.38
71	2,128	(b)	(a)	---	---	4,098	63.10
72	3,046	(b)	(a)	---	---	4,876	65.93
73	3,294	995	(a)	---	---	5,249	80.33
74	6,678	1,500	(a)	---	---	8,323	105.18
75	7,513	1,500	437	---	---	9,453	84.15
76	9,183	965	976	---	---	12,309	44.00
77	9,399	1,516	439	---	---	11,354	37.30
78	6,686	1,640	---	---	---	8,326	32.90
79	4,516	1,110	---	---	---	5,626	29.85
1880	2,439	422	---	---	---	2,561	31.00
81	2,194	---	---	---	---	2,194	29.83
82	2,171	---	---	---	---	2,171	28.23
83	1,894	---	---	---	---	1,894	28.75
84	881	---	(c)	---	---	881	30.50
85	385	---	---	---	---	385	30.75
86	409	---	---	---	---	409	35.60
87	673	---	---	---	---	673	42.38
88	126	---	---	---	---	126	42.50
89	812	---	---	---	---	812	45.00
1890	505	---	---	---	---	505	52.50
91	442	---	---	---	---	442	45.25
92	728	---	612	---	---	728	40.71
93	1,012	---	578	---	---	1,012	36.75
94	1,200	---	---	---	---	1,200	30.70
95	163	---	455	---	---	618	37.04
96	1,906	---	598	---	---	2,504	34.96
97	1,550	---	453	---	---	2,003	37.28
98	1,012	---	790	---	---	1,802	38.23
99	882	---	542	---	---	1,424	47.70
1900	816	---	---	---	(e)	816	44.94
01	1,545	---	400	---	---	1,945	48.46
02	1,950	---	317	---	---	2,267	43.20
03	1,730	---	284	---	---	2,014	42.25
04	913	---	399	---	---	1,312	37.62
05	223	---	250	---	---	473	35.94
06	---	---	---	---	---	---	36.50
07	100	---	---	---	---	100	38.16
08	100	---	---	---	---	100	42.33
09	17	---	---	---	---	17	47.71
1910	33	---	---	---	---	33	45.23
11	140	---	---	---	---	140	46.01
12	287	---	---	---	---	287	42.04
13	287	(d)	---	---	---	287	40.23
14	77	15	---	---	---	92	49.05
15	58	14	---	---	---	72	81.52

Year	Knoxville (Excelsior, Redington, Boston)	Reed (California, Andalusia)	Manhattan (Lake)	Red Elephant	Harrison (New England, Ruby, Grizzly, January)	Total ¹	Average price of quicksilver per flask
16	60	---	46	---	---	108	93.50
17	42	14	---	---	13	69	98.29
18	---	---	---	---	30	30	114.03
19	---	---	---	---	220	220	89.04
1920	---	---	---	---	---	---	75.45
21	---	---	---	---	---	---	44.56
22	68	---	---	---	---	68	55.35(f)
23	---	---	---	---	---	---	60.98
24	---	---	---	---	---	---	68.33
25	---	---	---	---	---	---	80.81
26	---	---	---	---	---	---	87.64
27	38	---	9	---	---	47	111.67
28	541	---	---	---	---	541	118.84
29	742	---	---	74	---	816	117.78
1930	970	---	---	154	---	1,124	110.36
31	649	5	---	90	---	744	83.22
32	13	2	1	---	---	16	52.30
33	21	---	3	---	---	24	55.94
34	74	---	---	15	---	89	67.22
35	38	---	---	16	---	54	67.23
36	12	---	2	---	2	16	76.62
37	---	---	131	---	28	159	90.18
38	---	---	72	---	28	100	75.47
39	---	4	33	---	14	51	103.94
1940	75	467	51	---	10	603	176.87
41	458	784	33	---	---	1,275	187.92
42	600	2,422	31	56	(d)	3,109	196.35
43	642	1,771	26	---	(d)	2,439	195.21
Total	121,967	15,146	11,562	405	345	156,458	

¹ 76.5 lbs. previous to June 1904; 75 pounds from June 1904 to January 1923; and 76 pounds since January 1928.

² Includes flasks of unknown origin before 1877.

(a) Manhattan production between 1863 and 1875 totaled 3,594 flasks.

(b) Amount of production unknown.

(c) Intermittent unrecorded production between 1884 and 1892.

(d) Small unrecorded production.

(e) Pre-1900 production small and unrecorded.

(f) Tariff Act of 1922 placed import duties of \$13.75 per flask.

Table 1—Production record of mines in Knoxville quicksilver district 1862-1943
(in flasks)²

the Clear Lake area, White and others (1973) concluded that the high-chloride connate waters found at Wilbur Springs are chemically and isotopically similar to the oil-field waters of the Central Valley of California, but are generally enriched in bicarbonates and boron. This water probably evolved from compaction and loss of pore water in the marine sediments, but is also affected to some extent by continuing low-grade metamorphism at depth.

Hydrocarbons present at Wilbur Springs and Knoxville are common in many of the mercury mines in this region and came to the surface in the same fractures that hot mineral laden water did. At Wilbur Springs, hot water and hydrocarbons (oil and gas) are ascending in the same fracture system.

Moisseeff (1966) concluded that the mercury at Wilbur Springs probably was concentrated from sediments in "solutions of metamorphic or connate origin." Although Barnes and others (1973) concluded that fluids presently discharged at Wilbur Springs could not have formed the quantity of mercury deposits there, since the concentration in the water is not high enough to account for the amount of mercury deposited since Tertiary time (?).

It is likely that the gold has a similar origin as the mercury. Steve Graham (personal communication, July 30, 1981), noting the lack of flour gold in gold-bearing gravels in the Sierras, sampled the Great Valley Sequence (and younger formations) in the spring of 1981, hoping to detect the timing of gold emplacement in the Sierras. He noted a general increase in gold with decrease in age. Knoxville Formation assayed .001-.002 oz. per ton, and the Eocene formations assayed .006-.008 oz. per ton. The high chloride content of the water may have facilitated in the transportation of gold, since in hydrothermal solutions in general gold is transported as the aurous chloride complex AuCl_2 (Radtke and Scheiner, 1970).

In the structurally complex Coast Range, several volcanic fields (Sonoma, Tolay, Berkely Hills, Quien Sabe, Neenach-Pinnacles) have erupted, generally decreasing in age northward. The Clear Lake volcanic field is the youngest and most northerly of these fields. Within the Clear Lake volcanic field, isotopic ages and the eruptive sequence also show a general decrease in age northward from 2 m.y. in the south to about 10,000 years in the north. The Geysers geothermal area is believed to be heated by a silicic magma body centered below 10 km in depth, the top of which is within 7 km of the surface (Hearn, and others 1981). On the basis of a detailed gravity study, a series of 100 m temperature gradient holes, and two deep wells (400 m and 1,200 m), Harrington and Verosub (1981) concluded that presumably the source of heat at Wilbur Springs is a deep magma intrusion. Fractures provide the porosity needed to transfer the heat in the form of hot water or steam from the intrusion.

The chemistry of mercury at the Wilbur Springs deposits has been studied in detail by Moisseeff (1966) and is beyond the scope of this paper. The geochemistry of the gold in these mines or the Knoxville mines is currently unknown. However, there are some conspicuous similarities between these deposits and the Carlin-type gold deposits which have become the source of large quantities of gold mined in Nevada.

Radtke and other workers (see Boyle, 1980, for a summary) have done the pioneer work on the Carlin-type deposits. From their work a checklist of geologic, mineralogic, and geochemical features common to these type of deposits has been developed. This list includes: silty carbonate rocks; the presence of hydrocarbons; gold associated with mercury, arsenic, antimony and thallium; replacement by fine-grained silica and minor pyrite, and; mineralization related with strong normal faults. Fluids at these deposits have been shown to be of meteoric origin. Work by Radtke and Scheiner (1970) suggest the presence of carbonaceous materials (hydrocarbons) may be critical for the deposition of gold.

In the Coast Ranges, mercury deposits are largely but not entirely, restricted to serpentine which has been replaced by silica and carbonate. Hydrocarbons and pyrite are very common in these deposits. The presence of the hydrocarbons supports the findings of White and others (1973) that the water which transported the mercury is of connate origin, and that the silty sediments of the Great Valley Sequence is the source. This water, therefore, has a different source than the Carlin-type deposits. Gold is associated with mercury and/or antimony in the Coast Ranges. Although on Red Mountain, southeast of Ukiah, gold is instead associated with copper. The mercury deposits of the Coast Ranges are late Tertiary to Quaternary in age. They are also associated with strong normal faults.

From this discussion, it can be seen that there are numerous similarities between the mercury deposits in the Coast Ranges of California and the Carlin-type gold deposits. The Knoxville gold deposit is the first major deposit of this type to be recognized in California. However, it is clear that there is potential for others.

MERCURY-GOLD DEPOSITS IN ADJACENT AREAS

In the Knoxville district, mercury deposits are confined to silica-carbonate rocks along northwest-trending faults, between the contact of serpentine and Knoxville sediments. The Manhattan mine is an exception, with mercury ore occurring in altered Tertiary olivine basalt and tuff. Minerals reported from these mines include cinnabar, metacinnabar, pyrite, marcasite, stibnite, native mercury, copper oxides, gold, millerite (nickel sulfide) and redingtonite (hydrous chromium sulfate). Pyrite in the district shows traces of gold (Averitt, 1945, p. 78). Apparently minor leaching has occurred, yielding free gold as reported by Becker (1888, p. 282):

The surface soil here (the Manhattan mine) and also near the Redington (Knoxville) mine contains cinnabar, resulting from the erosion of croppings, and accompanying the cinnabar is free gold, which may be found by panning the soil.

At Wilbur Springs, mercury ore occurs in the same localities as the hot springs. Three groups of mines can be recognized in the vicinity of Wilbur Springs. The Wilbur Springs Group consists of the Wide Awake, Central and Empire mines located along a "dike" of detrital serpentine, and the Manzanita and Cherry Hill mines located in fractured sedimentary

parties and other workers (see Fowler, 1935, for a summary) have done the
 present work on the Carlin-type deposits. From their work a checklist of
 geologic, stratigraphic, and geochemical features common to these types of
 deposits has been developed. This list includes: silty carbonate rocks;
 the presence of bedded argillaceous siltstone, argillaceous, and
 clay and limestone replacement of fine-grained siltstone and argillaceous
 sand, mineralization related with strong normal faults. Fluids at Carlin
 deposits have been shown to be of meteoric origin. Work by Rastin and
 Schaner (1970) suggest the presence of carbonate-sulfate brines (hydro-
 carbonates) may be critical for the deposition of gold.

In the Great Basin, many deposits are largely not entirely re-
 stricted to sedimentation which has been related by siltstone and carbonate.
 Hydrothermal and basaltic are very common in these deposits. The presence
 of the hydrothermal supports the findings of White and others (1970) that
 the water which transported the material is of carbonate origin, and that the
 silty sandstone of the Great Valley Sequence is the source. This water
 therefore, has a different source than the Carlin-type deposits. Gold is
 associated with many other minerals in the Great Basin. Although no
 Red Mountain, southeast of Union, gold is instead associated with copper.
 The mercury, arsenic of the Great Basin are also related to hydrothermal
 in origin. They are also associated with strong normal faults.

From this discussion, it can be seen that there are numerous similarities
 between the mercury deposits in the Great Basin of California and the
 Carlin-type gold deposits. The Nevada gold deposits is the first major
 deposit of this type to be recognized in California. However, it is clear
 that there is potential for others.

NEVADA-GOLD DEPOSITS IN ADJACENT AREAS

In the Nevada district, mercury deposits are confined to silt-
 stone and claystone along north-south trending faults between the contact of
 granitic and basaltic andesite. The Nevada area is an exception
 with mercury ore occurring in argillaceous siltstone and clay.
 Nevada deposits also have other minerals including malachite, pyrite,
 arsenic, antimony, native mercury, copper, silver, gold, arsenic
 (native and arsenic) and redoxymorphic (hydroxide and arsenic). Pyrite is
 the district where traces of gold (Averitt, 1935, p. 78). Presumably
 silver, including has occurred, including free gold as reported by Becker
 (1935, p. 282).

The surface gold belt (the Nevada area) and also near the
 Red Mountain (Nevada) area contains minerals, resulting from the
 erosion of granitic, and suggesting the minerals in these
 gold, which may be found by mining the soil.

At Union Springs, mercury and arsenic in the same localities as the gold
 deposits. These deposits are also can be recognized in the vicinity of
 Union Springs. The Union Springs area consists of the wide, low
 Great and Pacific ranges located along a "line" of central extension,
 and the mountains and Great Salt Lake located in the Great Basin.

rocks of the Knoxville Formation. The Elgin Group occurs in a body of silicified serpentine. The Clyde mine about a mile to the northwest, is at the contact of serpentine and the Knoxville formation. The Abbott Group includes the Boggess, Abbott and Turkey Run mines, which are located in silica-carbonate rock at the contact between the serpentine with Knoxville sediments. Minerals present in the Wilbur Springs district include cinnabar, metacinnabar, pyrite, marcasite, gold, and minor calomel.

Gold occurs at the Cherry Hill, Manzanita, Abbott, and Clyde mines (Logan, 1929). At the Abbott mine, traces of gold are present in the marcasite (Moisseff, 1966, p. 85). Becker (1888, p. 367) noted the ore at the Manzanita mine consisted of:

...cinnabar and gold, which are sometimes in direct contact, and some metacinnabarite. These minerals are accompanied by pyrite and marcasite, chalcopryite, stibnite, calcite, and quartz. The gold is often visible in feather-like, crystalline aggregates, sometimes in direct contact with cinnabar and sometimes deposited directly upon calcite....

At this mine, Moisseff (1966) described the occurrence of gold as specks or "sheets" in leached zones. The destruction of iron sulfides probably liberated the gold which then became scattered in the leached zones.

MERCURY-GOLD WITHIN THE WSA

The Shamrock mine situated in Sec. 23, 14T.13N., R.6W.MDM was located and worked prior to 1903 (Forstner, 1903). In 1927 and 1935, Orville Blevins of Redding produced mercury from the property. Foyle Mason, acquired the property in the early 1940s and held it until his death in 1980. Mason did not produce any mercury from the property. In 1968, M.C. Smith and the late Kay Miller, both of Redding, located the Deep Shaft and Zodiac claims over Mason's claims (the Shamrock and Merle) apparently with his permission. Smith and Miller erected a mill at the expense of \$100,000 and have produced a "couple of hundred" flasks of mercury. They determined the presence of gold on these claims, but were more interested in the mercury.

The Hunting Fault, which separates serpentine on the south from sediments of the Great Valley Sequence on the north, trends northwest across the Deep Shaft and Zodiac claims. The fault zone is expressed as a red clay gouge zone that contains sparse mercury values. The most significant mercury mineralization occurs in serpentine which is altered to opalite, located south of the fault. The altered serpentine is mineralized with cinnabar, copper oxides, pyrite, and gold. Small amounts of gold have been recovered from the ore by the use of a shaker table. Placer gold, and a nugget which weighed 0.25 ounces have been recovered from Rocky Creek. Reportedly, cinnabar crystals were found to have grown within one gold nugget (M.C. Smith personal communication, August 25, 1981).

In the southeastern end of the WSA, claims have been located in Sections 14, 15, 22 and 23 T.12N., R.6W. and Sections 7, 18 and 19 T.12N., R.5W.MDM. These claims were located on the periphery of the recently discovered immense gold deposit at Knoxville.

CHROMITE

Placer chromite occurs east of Deadman Canyon in Sections 7 and 13 T.13N., R.6W. (Brice, 1953).

ABESTOS

Asbestos has been prospected north of Brushy Sky High in the SE 1/4 of Section 34 T.13N., R.6W.MDM. Here, a chrysotile vein trends N.44°E. and dips 52°W in highly sheared serpentine. The vein is 18 inches wide at the point of discovery and pinches to 6 inches within 100 feet along strike. The asbestos fibers average an eighth of an inch long and are slightly brittle. In 1952, soil was removed with a bulldozer and prospect trenches were cut at four points across the vein (Brice, 1953 p. 60).

GEMSTONES

Jasper cobbles are common along Cache Creek (Brice, 1953).

MAGNESITE

Magnesite of probably minor quantities occurs at the Shamrock mercury mine (Foyle Mason mining claim file CAMC 49164).

GRAVEL

Gravel deposits along the north fork of Cache Creek from Upper Wilson Valley to the confluence with Long Valley Creek have potential for use in concrete aggregate (Klein and Goldman, 1958).

OIL AND GAS

Five oil seepages in the Wilbur Springs area have prompted the drilling of five shallow wells on the Wilbur Springs Anticline between 1844 and 1937. All of the wells were less than 3,000 feet deep. None of these wells has resulted in commercial production, but there

In the southeastern end of the area, claims have been located in sections 14, 15, 16 and 17, T.13N., R.10W., and sections 18 and 19, T.13N., R.10W. These claims were located on the north side of the recently discovered massive gold deposit at Rockville.

ROCKVILLE

Massive deposits occur east of Garden Canyon in sections 18 and 19, T.13N., R.10W. (Pattee, 1933).

ROCKVILLE

Section has been prospected north of Quarry 25, high in the SE 1/4 of section 34, T.13N., R.10W. Here, a chrysotile vein trends N. 40° E. and dips 35° N. in highly altered sedimentary. The vein is 18 inches wide at the point of discovery and narrows to 6 inches within 100 feet along strike. The section there shows an outcrop of an iron ore and was slightly drilled. In 1933, drill was removed with a bulldozer and exposed strata were cut at four points across the vein (Pattee, 1933 p. 60).

ROCKVILLE

Large deposits are common along Cedar Creek (Pattee, 1933).

ROCKVILLE

Deposits of probably silver sulfides occur at the Canyon NW-1, only one (open) being shown (Pattee, 1933).

ROCKVILLE

Small deposits along the north fork of Cedar Creek from Canyon NW-1 to the south side with Long Valley Creek have been shown. This is now in completely igneous (Pattee and Coleman, 1930).

ROCKVILLE

Five old adits in the Willow Springs area have produced the following quantities of silver: Willow Springs No. 1, 1,000 lbs. silver; Willow Springs No. 2, 1,000 lbs. silver; Willow Springs No. 3, 1,000 lbs. silver; Willow Springs No. 4, 1,000 lbs. silver; Willow Springs No. 5, 1,000 lbs. silver. All of the silver was less than 1,000 lbs. each.

have been shows of oil in several of them (Lawton, 1956, p. 211). At the Reed Mine, north of Knoxville, Averitt (1945, p. 87) observed oil seeps.

Within the Cache Creek area gas seeps were observed by Lawton (1956, p. 194):

...where the Wilson fault crosses Cache Creek just south of Section 21, T.13., R.5W. Sulfurous springs are also common in this area.

Lawton (op. cit., p. 214) considered the Cache Creek area one of the most promising in the area of his thesis for the entrapment of oil. He stated:

In the southcentral portion of the area, in parts of Section 16, 17, 18, 20 and 21, T.13N., R.5W., the gentle folding and the Dunfield, Wilson and Kennedy Faults may form a structural trap. Drilling objectives here would be the Leesville member of the Little Valley formation, the Blue Ridge member of the Crack Canyon Formation, and the sandstone and conglomerate horizons in the Sulfur Creek (=? Knoxville) Formation. This is one of the most promising structures in this area as it is large and the bounding faults and the gentle folding are well defined. The best well location might be in the center of Section 21, T.13N., R.5W.

GEO THERMAL ENERGY

The Wilbur Springs geothermal area has the potential to generate electric power (Higgins and Martin, 1980). A series of shallow temperature gradient holes indicate thermal gradients as high as $.3^{\circ}\text{C}/\text{m}$, and two deep holes drilled to 400 m and 1,200 m reached maximum bottom temperatures of 120°C and 140°C respectively. Harrington and Verosub (1981) studied the Wilbur Springs area, and concluded that the geothermal reservoir, supplying heat for the hot springs continues south of Highway 20 in the vicinity of Destanella Flat. This is about one mile from the WSA.

Very little is known about the geothermal potential of the Knoxville KGRA. There is a 22°C warm spring at Knoxville.

have been shown to all in several of them (Lawton, 1956, p. 211).
At the last mine, north of Knoxville, Avestic (1957, p. 87) observed
oil seeps.

Within the Lenoir Creek area gas seeps were observed by Lawton (1956,
p. 194):

...where the Wilson fault crosses Lenoir Creek just south of Sec-
tion 21, T.15, R.2W. Sulfur gas seeps are also common in this
area.

Lawton (loc. cit., p. 194) considered the Lenoir Creek area one of the most
promising in the area of his thesis for the entrapment of oil. He stated:

In the southeastern portion of the area, in parts of Section 16,
17, 18, 20 and 21, T.15N, R.2W, the gentle folding and the
unusually high permeability faults may form a structural trap.
Drilling objectives here would be the Knoxville member of the
Little Valley formation, the Blue Ridge member of the Creek
Group formation, and the sandstone and conglomerate formations in
the Lenoir Creek (or Knoxville) formation. This is one of the
most promising structures in this area as it is large and the
favorable faults and the gentle folding are well defined. The
best well location might be in the center of Section 21, T.15N,
R.2W.

GEOLOGICAL SUMMARY

The Wilson Springs geothermal area has the potential to generate electricity
power (Hodge and Weller, 1980). A series of shallow temperature gradi-
ent holes indicate thermal gradients as high as 150°C/m, and two deep holes
drilled to 500 m and 1,300 m reached maximum bottom temperatures of 120°C
and 140°C respectively. Harrington and Weller (1981) studied the Wilson
Springs area and concluded that the geothermal reservoir, supplying heat
for the hot springs, extends south of Highway 20 in the vicinity of
Knoxville Flat. This is about one mile from the WEA

Very little is known about the geothermal potential of the Knoxville
area. There is a 1961 well being drilled at Knoxville.

Bibliography

- Averitt, P., 1945. Quicksilver deposits of the Knoxville district Napa, Yolo and Lake Counties: California Journal of Mines and Geology, V. 41, No. 2.
- Bailey, E.H., Blake, M.C. Jr., Jones, D.L., 1970. On land mesozoic oceanic crust in California Coast Ranges: U.S. Geological Survey Professional Paper 700-C, pp. C70-C81.
- Bailey, E.H., Irwin, W.P., Jones, D.L., 1964. Franciscan and related rocks, and their significance in the geology of Western California: California Division of Mines and Geology Bulletin 183.
- Bailey, E.H., Meyers, W.B., 1949. Quicksilver and antimony deposits of the Stayton District, California: in California Division of Mines Bulletin 147, pp. 37-56.
- Barnes, I., and others, 1973. Chemical composition of naturally occurring fluids in relation to mercury deposits in part of North-Central California: U.S. Geological Survey Bulletin 1382-A.
- Becker, G.F., 1888. Geology of the quicksilver deposits of the Pacific Slope: U.S. Geological Survey Monograph 13.
- Boyle, R.W., 1980. The geochemistry of gold and its deposits: Canada Geological Survey Bulletin 280.
- Bradley, W.W., 1916. The Counties of Colusa, Glenn, Lake, Marin, Napa, Solano, Sonoma, Yolo: California Mining Bureau Report 14.
- Brice, J.C., 1953. Geology of Lower Lake quadrangle California: California Division of Mines Bulletin 166.
- Bufe, C.G. and others, 1981. Seismicity of the Geysers-Clear Lake Region: U.S. Geological Survey Professional Paper 1141, pp. 129-317.
- Carlson, C., 1981. Sedimentary serpentinites of the Wilbur Springs quadrangle - a possible early Cretaceous structural and stratigraphic link between the Franciscan Complex and the Great Valley Sequence: Stanford University, MS Thesis.
- Carlson, C.R., 1962. Oil seeps and early petroleum development in Northern California: California Division of Mines Geology Bulletin 181.
- Casteel, R.W., and Rymer, M.J., 1981. Pliocene and Pleistocene fishes from the Clear Lake area: U.S. Geological Survey Professional Paper 1141, pp. 231-235.
- Donnelly-Nolan, J.M., and others, 1981. Geochronology and evolution of the Clear Lake Volcanics: U.S. Geologic Survey Professional Paper 1141, pp. 47-60.

- Donnelly-Nolan, J.M., McLaughlin, R.J., Hearn, B.C., Jr., 1980. Tectonic influences on hydrothermal and magmatic systems in the Geysers-Clear Lake Region, CA.: Transactions American Geophysical Union Abstracts, Vol. 61, No. 46, pp. 1149-1150.
- Forstner, W., 1903. The quicksilver resources of California: California Mining Bureau Bulletin 27.
- Harrington, J.M. and Verosub, K.L., 1981. A detailed gravity survey of the Wilbur Springs area, California: U.S. Geological Survey Professional Paper 1141, pp. 161-166.
- Hearn, B.C., and others, 1981. The Clear Lake Volcanics: tectonic setting and magma sources: U.S. Geologic Survey Professional Paper 1141, pp. 25-45.
- Homestake Mining Company: Press Release, May 12, 1981.
- Klein, I.E., Goldman, H.B., 1958. Sand and gravel resources of Cache Creek in Lake, Colusa, and Yolo Counties, California: California Journal of Mines and Geology Vol. 54, No. 1.
- Lawton, J.E., 1956. Geology of the north half of the Morgan Valley quadrangle and the south half of the Wilbur Springs quadrangle: Stanford University, PhD Thesis.
- Logan, C.A., 1929. Sacramento Field Division, Colusa County: Mining in California, Vol. 25, No. 3.
- McLaughlin, R.J., 1981. Tectonic setting of pre-tertiary rocks and its relation to geothermal resources in the Geysers-Clear Lake Area: U.S. Geologic Survey Professional Paper 1141, pp. 3-21.
- McLaughlin, R.J., Donnelly-Nolan, J.M. (editors), 1981. Research in the Geysers-Clear Lake geothermal area, Northern California: U.S. Geological Survey Professional Paper 1141.
- Metallurgical testing of Oat Hill mercury areas: Newmont Exploration Limited, File 222-65-1, September 22, 1965: Unpublished Manuscript, California Room, California State Library
- Moisseeff, A.N., 1966. The geology and geochemistry of the Wilbur Springs quicksilver district, Colusa and Lake Counties, California: Stanford University, PhD Thesis.
- Page, B.M., 1966. Geology of the coast ranges of California: California Division of Mines and Geology Bulletin 190, pp. 255-275.
- Radtko, A.S., Rye, R.O., Dickson, F.W., 1980. Geology and stable isotope studies of the Carline gold deposit Nevada: Economic Geology Vol. 75, No. 5, pp. 641-672.

Wells, J. H., 1960. Tectonic
influence on hydrothermal and magmatic systems in the Cascade-Lake
Lake Region, California: Transactions American Geophysical Union, Washington,
Vol. 41, no. 46, pp. 1199-1200.

Wells, J. H., 1961. The geothermal resources of California: California
Mining Bureau Bulletin 27.

Wells, J. H. and Wright, K. L., 1961. A detailed gravity survey of the
White Sulphur area, California: U.S. Geological Survey Professional
Paper 1141, pp. 141-166.

Wells, J. H., and others, 1961. The Clear Lake Volcanics: tectonic setting
and geologic structure: U.S. Geological Survey Professional Paper 1141, pp.
23-43.

Wells, J. H., 1961. Press Release, May 12, 1961.

Wells, J. H., 1961. Some new gravity resources of Lake County,
California, and Lake County, California: California Journal of
Mining and Geology Vol. 41, no. 1.

Wells, J. H., 1961. Geology of the north half of the Upper Valley and
range and the south half of the White Sulphur Mountains: Stanford
University, PhD thesis.

Wells, J. H., 1961. Geomorphological Field Division, California Energy
Commission, Vol. 1, no. 1.

Wells, J. H., 1961. Tectonic setting of pre-tertiary rocks and its im-
pact on geothermal resources in the Cascade-Lake Lake Area: U.S.
Geological Survey Professional Paper 1141, pp. 3-21.

Wells, J. H., 1961. Research in the
Lake County-Lake County area, Northern California: U.S. Geo-
logical Survey Professional Paper 1141.

Wells, J. H., 1961. Geology of the Lake County-Lake County area, Northern California:
U.S. Geological Survey Professional Paper 1141, pp. 22-23.

Wells, J. H., 1961. The geology and geothermal resources of the White Sulphur
Mountains: California Journal of Mining and Geology Vol. 41, no. 1.

Wells, J. H., 1961. Geology of the Lake County-Lake County area, Northern California:
U.S. Geological Survey Professional Paper 1141, pp. 24-25.

Wells, J. H., 1961. Geology and geothermal resources of the Lake County-Lake County area, Northern California:
U.S. Geological Survey Professional Paper 1141, pp. 26-27.

Radtke, A.S., Scheiner, B.J., 1970. Carlin gold deposit, Nevada: The role of carbonaceous materials in gold deposition: Economic Geology Vol. 65, No. 2, pp. 87-102.

Real, C.R., Toppozada, T.R., Parke, D.L., 1978. Earthquake Epicenter Map of California: California Division of Mines and Geology Map Sheet 39.

White, D.E., Barnes, I., O'Neil, J.R., 1973. Thermal and mineral waters of nonmeteoric origin, California Coast Ranges: Geological Society of America Bulletin V.84, pp. 597-560.

Wagner, A.E., Stalder, B.J., 1970. Early gold deposits, Nevada: The role
of subvolcanic materials in gold deposition. Economic Geology, vol.
65, no. 2, pp. 27-102.

Reel, D.E., Johnson, R.E., Paine, D.L., 1978. Earthquake epicenter map
of California. California Division of Mines and Geology Map Sheet 29.

Miller, D.E., Barnes, J., Goff, J.R., 1972. Thermal and mineral waters of
northeastern Oregon. California Coast Range. Geological Society of
America Bulletin 83, pp. 205-222.

BLM Library
Bldg. 50
Denver Federal Center
P.O. Box 25047
Denver, Colorado 80225